professionals that treat	pharmacist, dentist and you. I am under antico ated tablets to prevent	agulation treatment with
Please complete this se	ction or ask your doctor	r to do it
Name:	Birth Date:	
Indication:	Dose:	mg twice daily
Doctor's Name:	Doctor's telepho	ne:
remember and continue	y as instructed. If you miss to follow your dosing sche aban without talking to you	a dose, take it as soon as you edule. ur doctor, as you are at risk of

- Apixaban helps to thin your blood. However, this may increase your risk of bleeding.
- Signs and symptoms of bleeding include bruising or bleeding under the skin, tar-coloured stools, blood in urine, nose-bleed, dizziness, tiredness, paleness or
- weakness, sudden severe headache, coughing up blood or vomiting blood. If the bleeding does not stop on its own, **seek medical attention immediately.**  If you need surgery or any invasive procedure, inform your doctor that
- you are taking Apixaban.

- Information for healthcare professionals
  Apixaban is an oral anticoagulant acting by direct selective inhibition of factor Xa.
  Apixaban may increase the risk of bleeding. In case of major bleeding
- events, it should be stopped immediately.Treatment with Apixaban does not require routine monitoring of exposure. A calibrated quantitative anti-Factor Xa assay may be useful in exceptional situations, e.g., overdose and emergency surgery (prothrombin time (PT), international normalised ratio (INR) and activated partial thromboplastin time
- (aPTT) clotting tests are not recommended) see SmPC. An agent to reverse the anti-factor Xa activity of apixaban is available. Date of preparation: June 2022

216234