

Package leaflet: Information for the user

Rivaroxaban 2.5 mg film-coated tablets

rivaroxaban

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Rivaroxaban is and what it is used for
2. What you need to know before you take Rivaroxaban
3. How to take Rivaroxaban
4. Possible side effects
5. How to store Rivaroxaban
6. Contents of the pack and other information

1. What Rivaroxaban is and what it is used for

You have been given Rivaroxaban because

- you have been diagnosed with an acute coronary syndrome (a group of conditions that includes heart attack and unstable angina, a severe type of chest pain) and have been shown to have had an increase in certain cardiac blood tests.

Rivaroxaban reduces the risk in adults of having another heart attack or reduces the risk of dying from a disease related to your heart or your blood vessels.

Rivaroxaban will not be given to you on its own. Your doctor will also tell you to take either:

- acetylsalicylic acid or
- acetylsalicylic acid plus clopidogrel or ticlopidine.

or

- you have been diagnosed with a high risk of getting a blood clot due to a coronary artery disease or peripheral artery disease which causes symptoms.

Rivaroxaban reduces the risk in adults of getting blot clots (atherothrombotic events).

Rivaroxaban will not be given to you on its own. Your doctor will also tell you to take acetylsalicylic acid.

- In some cases, if you get Rivaroxaban after a procedure to open a narrowed or closed artery of your leg to restore blood flow, your doctor may also prescribe clopidogrel for you to take in addition to acetylsalicylic acid for a short while.

Rivaroxaban contains the active substance rivaroxaban and belongs to a group of medicines called antithrombotic agents. It works by blocking a blood clotting factor (factor Xa) and thus reducing the tendency of the blood to form clots.

2. What you need to know before you take Rivaroxaban

Do not take Rivaroxaban

- **if you are allergic** to rivaroxaban or any of the other ingredients of this medicine (listed in section 6)
 - **if you are bleeding excessively**
 - **if you have a disease or condition in an organ of the body that increases the risk of serious bleeding (e.g. stomach ulcer, injury or bleeding in the brain, recent surgery of the brain or eyes)**
 - if you are taking medicines to prevent blood clotting (e.g. warfarin, dabigatran, apixaban or heparin), except when changing anticoagulant treatment or while getting heparin through a venous or arterial line to keep it open
 - **if you have an acute coronary syndrome and previously had a bleeding or a blood clot in your brain (stroke)**
 - **if you have** coronary artery disease or peripheral artery disease and previously had a bleeding in your brain (stroke) or where there was a blockage of the small arteries providing blood to the brain's deep tissues (lacunar stroke) or if you had a blood clot in your brain (ischaemic, non-lacunar stroke) in the previous month
 - **if you have a liver disease** which leads to an increased risk of bleeding
 - **if you are pregnant or breast-feeding**
- Do not take Rivaroxaban and tell your doctor** if any of these apply to you.

Warnings and precautions

Talk to your doctor or pharmacist before taking Rivaroxaban.

Rivaroxaban should not be used in combination with certain other medicines which reduce blood clotting such as prasugrel or ticagrelor other than acetylsalicylic acid and clopidogrel/ticlopidine.

Take special care with Rivaroxaban

- if you have **an increased risk of bleeding, as could be the case in situations** such as:
 - **severe kidney disease, since your kidney function may affect the amount of medicine that works in your body**
 - **if you are taking other medicines to prevent blood clotting (e.g. warfarin, dabigatran, apixaban or heparin), when changing anticoagulant treatment or while getting heparin through a venous or arterial line to keep it open (see section “Other medicines and Rivaroxaban”)**
 - **bleeding disorders**
 - **very high blood pressure, not controlled by medical treatment**
 - **diseases of your stomach or bowel that might result in bleeding, e.g. inflammation of the bowels or stomach, or inflammation of the oesophagus (gullet), e.g. due to gastroesophageal reflux disease (disease where stomach acid goes upwards into the oesophagus) or tumours located in the stomach or bowels or genital tract or urinary tract**
 - **a problem with the blood vessels in the back of your eyes (retinopathy)**
 - a lung disease where your bronchi are widened and filled with pus (bronchiectasis), or previous bleeding from your lung
 - you are older than 75 years
 - you weigh 60 kg or less
 - you have a coronary artery disease with severe symptomatic heart failure
- if you have a prosthetic heart valve
- if you know that you have a disease called antiphospholipid syndrome (a disorder of the immune system that causes an increased risk of blood clots), tell your doctor who will decide if the treatment may need to be changed.

If any of the above apply to you, tell your doctor before you take Rivaroxaban. Your doctor will decide, if you should be treated with this medicine and if you should be kept under closer observation.

If you need to have an operation

- it is very important to take Rivaroxaban before and after the operation exactly at the times you have been told by your doctor.

- If your operation involves a catheter or injection into your spinal column (e.g. for epidural or spinal anaesthesia or pain reduction):
 - it is very important to take Rivaroxaban before and after the injection or removal of the catheter exactly at the times you have been told by your doctor

- tell your doctor immediately if you get numbness or weakness of your legs or problems with your bowel or bladder after the end of anaesthesia, because urgent care is necessary.

Children and adolescents

Rivaroxaban is **not recommended for people under 18 years of age**. There is not enough information on its use in children and adolescents.

Other medicines and Rivaroxaban

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

– **If you are taking**

- some **medicines for fungal infections** (e.g. fluconazole, itraconazole, voriconazole, posaconazole), unless they are only applied to the skin
- ketoconazole tablets (used to treat Cushing’s syndrome - when the body produces an excess of cortisol)
- some medicines for bacterial infections (e.g. clarithromycin, erythromycin)
- some **anti-viral medicines for HIV / AIDS** (e.g. ritonavir)
- other medicines to **reduce blood clotting** (e.g. enoxaparin, clopidogrel or vitamin K antagonists such as warfarin and acenocoumarol, prasugrel and ticagrelor (see section “Warnings and Precautions”))
- **anti-inflammatory and pain relieving medicines** (e.g. naproxen or acetylsalicylic acid)
- dronedarone, a medicine to treat abnormal heart beat
- some medicines to treat depression (selective serotonin reuptake inhibitors (SSRIs) or serotonin norepinephrine reuptake inhibitors (SNRIs))

If any of the above apply to you, tell your doctor before taking Rivaroxaban, because the effect of Rivaroxaban may be increased. Your doctor will decide, if you should be treated with this medicine and if you should be kept under closer observation.

If your doctor thinks that you are at increased risk of developing stomach or bowel ulcers, he may also use a preventative ulcer treatment.

– If you are taking

- some medicines for treatment of epilepsy (phenytoin, carbamazepine, phenobarbital)
- St John’s Wort (*Hypericum perforatum*), a herbal product used for depression
- rifampicin, an antibiotic

If any of the above apply to you, tell your doctor before taking Rivaroxaban, because the effect of Rivaroxaban may be reduced. Your doctor will decide, if you should be treated with Rivaroxaban and if you should be kept under closer observation.

Pregnancy and breast-feeding

Do not take Rivaroxaban if you are pregnant or breast-feeding. If there is a chance that you could become pregnant, use a reliable contraceptive while you are taking Rivaroxaban. If you become pregnant while you are taking this medicine, tell your doctor immediately, who will decide how you should be treated.

Driving and using machines

Rivaroxaban may cause dizziness (common side effect) or fainting (uncommon side effect) (see section 4, “Possible side effects”). You should not drive, ride a bicycle or use any tools or machines if you are affected by these symptoms.

Rivaroxaban contains lactose and sodium

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially “sodium-free”.

3. How to take Rivaroxaban

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

How much to take

The recommended dose is one 2.5 mg tablet twice a day. Take Rivaroxaban around the same time every day (for example, one tablet in the morning and one in the evening). This medicine can be taken with or without food.

If you have difficulty swallowing the tablet whole, talk to your doctor about other ways to take Rivaroxaban. The tablet may be crushed and mixed with water or apple puree immediately before you take it.

If necessary, your doctor may also give you the crushed Rivaroxaban tablet through a stomach tube.

Rivaroxaban will not be given to you on its own.

Your doctor will also tell you to take acetylsalicylic acid. If you get Rivaroxaban after an acute coronary syndrome, your doctor may tell you to also take clopidogrel or ticlopidine.

If you get Rivaroxaban after a procedure to open a narrowed or closed artery of your leg to restore blood flow, your doctor may also prescribe clopidogrel for you to take in addition to acetylsalicylic acid for a short while.

Your doctor will tell you how much of these to take (usually between 75 to 100 mg acetylsalicylic acid daily or a daily dose of 75 to 100 mg acetylsalicylic acid plus a daily dose of either 75 mg clopidogrel or a standard daily dose of ticlopidine).

When to start Rivaroxaban

Treatment with Rivaroxaban after an acute coronary syndrome should be started as soon as possible after stabilisation of the acute coronary syndrome, at the earliest 24 hours after admission to hospital and at the time when parenteral (via injection) anticoagulation therapy would normally be stopped.

Your doctor will tell you when to start treatment with Rivaroxaban if you have been diagnosed with coronary artery disease or peripheral artery disease.

Your doctor will decide how long you must continue treatment.

If you take more Rivaroxaban than you should

Contact your doctor immediately if you have taken too many Rivaroxaban tablets. Taking too much Rivaroxaban increases the risk of bleeding.

If you forget to take Rivaroxaban

Do not take a double dose to make up for a missed dose. If you miss a dose, take your next dose at the usual time.

If you stop taking Rivaroxaban

Take Rivaroxaban on a regular basis and for as long as your doctor keeps prescribing it.

Do not stop taking Rivaroxaban without talking to your doctor first. If you stop taking this medicine, it may increase your risk of having another heart attack or stroke or dying from a disease related to your heart or your blood vessels.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Like other similar medicines to reduce the formation of blood clots, Rivaroxaban may cause bleeding which may potentially be life threatening. Excessive bleeding may lead to a sudden drop in blood pressure (shock). In some cases the bleeding may not be obvious.

Tell your doctor immediately if you experience any of the following side effects:

• **Signs of bleeding**

- bleeding into the brain or inside the skull (symptoms can include headache, one-sided weakness, vomiting, seizures, decreased level of consciousness, and neck stiffness. A serious medical emergency. Seek medical attention immediately!)
- long or excessive bleeding
- exceptional weakness, tiredness, paleness, dizziness, headache, unexplained swelling, breathlessness, chest pain or angina pectoris.

Your doctor may decide to keep you under closer observation or change the treatment.

Signs of severe skin reactions

- spreading intense skin rash, blisters or mucosal lesions, e.g. in the mouth or eyes (Stevens-Johnson syndrome/toxic epidermal necrolysis)..
- a drug reaction that causes rash, fever, inflammation of internal organs, blood abnormalities and systemic illness (DRESS syndrome). The frequency of these side effects is very rare (up to 1 in 10,000 people).

Signs of severe allergic reactions

- swelling of the face, lips, mouth, tongue or throat; difficulty swallowing; hives and breathing difficulties; sudden drop in blood pressure. The frequencies of severe allergic reactions are very rare (anaphylactic reactions, including anaphylactic shock; may affect up to 1 in 10,000 people) and uncommon (angioedema and allergic oedema; may affect up to 1 in 100 people).

Overall list of possible side effects

Common (may affect up to 1 in 10 people)

- reduction in red blood cells which can make the skin pale and cause weakness or breathlessness
- bleeding in the stomach or bowel, urogenital bleeding (including blood in the urine and heavy menstrual bleeding), nose bleed, bleeding in the gum
- bleeding into the eye (including bleeding from the whites of the eyes)
- bleeding into tissue or a cavity of the body (haematoma, bruising)
- coughing up blood
- bleeding from the skin or under the skin
- bleeding following an operation
- oozing of blood or fluid from surgical wound
- swelling in the limbs
- pain in the limbs
- impaired function of the kidneys (may be seen in tests performed by your doctor)
- fever
- stomach ache, indigestion, feeling or being sick, constipation, diarrhoea
- low blood pressure (symptoms may be feeling dizzy or fainting when standing up)
- decreased general strength and energy (weakness, tiredness), headache, dizziness
- rash, itchy skin
- blood tests may show an increase in some liver enzymes

Uncommon (may affect up to 1 in 100 people)

- bleeding into the brain or inside the skull (see above, signs of bleeding)
- bleeding into a joint causing pain and swelling
- thrombocytopenia (low number of platelets, which are cells that help blood to clot)
- allergic reactions, including allergic skin reactions
- impaired function of the liver (may be seen in tests performed by your doctor)
- blood tests may show an increase in bilirubin, some pancreatic or liver enzymes or in the number of platelets
- fainting
- feeling unwell
- faster heartbeat
- dry mouth
- hives

Rare (may affect up to 1 in 1,000 people)

- bleeding into a muscle
- cholestasis (decreased bile flow), hepatitis incl. hepatocellular injury (inflamed liver incl. liver injury)
- yellowing of the skin and eye (jaundice)
- localised swelling
- collection of blood (haematoma) in the groin as a complication of the cardiac procedure where a catheter is inserted in your leg artery (pseudoaneurysm)

Not known (frequency cannot be estimated from the available data)

- kidney failure after a severe bleeding
- increased pressure within muscles of the legs or arms after a bleeding, which leads to pain, swelling, altered sensation, numbness or paralysis (compartment syndrome after a bleeding)

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.co.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Rivaroxaban

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and on each blister or bottle after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Rivaroxaban contains

- The active substance is rivaroxaban. Each tablet contains 2.5 mg of rivaroxaban.
- The other ingredients are:
Tablet core: cellulose, microcrystalline, lactose monohydrate, sodium laurilsulphate, hypromellose, croscarmellose sodium, magnesium stearate. See section 2 “Rivaroxaban contains lactose and sodium”.

Tablet film coat: hypromellose, titanium dioxide, macrogol, iron oxide yellow.

What Rivaroxaban looks like and contents of the pack

Rivaroxaban 2.5 mg film-coated tablets are yellow, round, biconvex (6 mm diameter), engraved with “2.5” on one side and plain on the other side.

They come:

- in blisters in cartons of 20, 28, 45, 56 or 196 film-coated tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Glenmark Pharmaceuticals Europe Limited

Laxmi House, 2-B Draycott Avenue, Kenton, Middlesex, HA3 0BU, United Kingdom

Manufacturer

S.C. Labormed-Pharma S.A.

Bd. Theodor Pallady nr. 44B, sector 3

Bucharest cod 032266

Romania

Adalvo Ltd

Malta Life Sciences Park,

Building 1, Level 4, Sir Temi Zammit Buildings, San Gwann

SGN 3000, Malta

Glenmark Pharmaceuticals s.r.o.

Fibichova 143

566 17 Vysoké Mýto

Czech Republic

Saneca Pharmaceuticals a.s.

Nitrianska 100,

920 27 Hlohovec,

Slovak Republic

This leaflet was last revised in 07/2023.