

## **Package leaflet: Information for the user**

### **Spironolactone Waymade 25 mg/5ml oral solution** spironolactone

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, or pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Spironolactone Waymade is and what it is used for
2. What you need to know before you take Spironolactone Waymade
3. How to take Spironolactone Waymade
4. Possible side effects
5. How to store Spironolactone Waymade
6. Contents of the pack and other information

#### **1. What Spironolactone Waymade is and what it is used for**

Spironolactone Waymade contains the active substance spironolactone. Spironolactone Waymade belongs to a group of medicines called ‘diuretics’ – you may know these as ‘water’ tablets. You may have gone to your doctor because you had swollen ankles or were short of breath. This can happen when your heart’s pumping action has become weak because of too much fluid in your body. This is called ‘congestive heart failure’. Pushing extra fluid around your body means your heart has to work harder. Your doctor has given you Spironolactone Waymade to help you lose the extra fluid from your body. This will mean your heart has to do less work. You lose the extra fluid as urine, so you may need to go to the toilet more often while you are taking Spironolactone Waymade.

You can also take Spironolactone Waymade for the following illnesses:

- ‘Nephrotic syndrome’ - a kidney disorder that causes too much fluid in your body
- ‘Ascites’ - too much fluid in your abdomen and ‘oedema’ - accumulation of fluid beneath skin or in one or more cavities of the body that produces swelling, for example caused by cirrhosis of the liver
- ‘Malignant ascites’ - fluid containing cancer cells that collect in the abdomen
- ‘Primary aldosteronism’ - extra fluid in your body caused by too much of a hormone called ‘aldosterone’.

If you have these illnesses, Spironolactone Waymade will help your body to get rid of the extra fluid. You must talk to a doctor if you do not feel better or if you feel worse.

Children should only be treated under guidance of a paediatric specialist.

#### **2. What you need to know before you take Spironolactone Waymade**

##### **Do not take Spironolactone Waymade if**

- you are allergic to spironolactone or any of the other ingredients of this medicine (listed in section 6)
- you cannot pass urine
- you have severe kidney disease

- you have Addison's disease; (a hormone deficiency characterised by extreme weakness, loss of weight and low blood pressure)
- you have hyperkalaemia (raised blood potassium levels)
- you are breast-feeding
- you are taking water tablets (potassium sparing diuretics) or any potassium supplements
- you are taking eplerenone (a medicine for high blood pressure).

Children with moderate to severe kidney disease must not take Spironolactone Waymade.

### **Warnings and precautions**

Talk to your doctor or, pharmacist or nurse before taking Spironolactone Waymade if

- you suffer from kidney disease especially children with hypertension or liver disease. Your doctor will routinely assess you particularly if you are elderly
- you have difficulty passing urine
- you have a disease that can result in electrolyte balance disturbance in your blood such as potassium or sodium
- you have severe heart failure
- you are pregnant

If you experience reduced kidney function or kidney failure you may have severe increases in the levels of potassium in your blood. This can affect the way your heart functions and in extreme cases this can be fatal.

Concomitant administration of Spironolactone Waymade with certain medicines, potassium supplements and food rich in potassium may lead to severe hyperkalaemia (increased potassium blood level). The symptoms of severe hyperkalaemia might include muscle cramps, irregular heart rhythm, diarrhoea, nausea, dizziness or headache.

### **Other medicines and Spironolactone Waymade**

Tell your doctor or pharmacist if you are taking, have recently taken or might take use any other medicines. Your doctor may wish to alter your dose of Spironolactone Waymade if you are taking any of the following:

- digoxin or carbenoxolone
- medicines for high blood pressure including angiotensin converting enzyme (ACE) inhibitors
- other diuretics
- non-steroidal anti-inflammatory drugs (NSAIDs) such as aspirin, indomethacin, mefenamic acid or ibuprofen
- potassium supplements
- heparin or low molecular weight heparin (medicines used to prevent blood clots)
- antipyrine
- medicines known to cause hyperkalaemia (raised blood potassium levels)
- trimethoprim and trimethoprim-sulfamethoxazole

Spironolactone Waymade reduces your responsiveness to noradrenaline. If you are going to have an operation where you will be given an anaesthetic, tell the doctor in charge that you are taking Spironolactone Waymade.

### **Tell your doctor, if you are using abiraterone for treatment of prostate cancer.**

### **Spironolactone Waymade with food, drink and alcohol**

See section 3 'How to take Spironolactone Waymade'.

### **Pregnancy, breast-feeding and fertility**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy

There are limited data from the use of spironolactone in pregnant women. Your doctor will only prescribe Spironolactone Waymade if the potential benefit outweighs the potential risk.

#### **Breast-feeding**

Spironolactone Waymade should not be used if you are breastfeeding. You should discuss the use of Spironolactone Waymade with your doctor, who will advise you to consider an alternative method of feeding your baby while you are taking this medicine.

#### **Driving and using machines**

Take care if you drive or operate machinery. Drowsiness and dizziness have been associated with Spironolactone Waymade treatment and this may affect your ability to drive or operate machinery safely.

### **3. How to take Spironolactone Waymade**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure. The volume of this medicine you need to take depends on your illness.

#### **Recommended dose**

This medicine should be taken once a day with food.

#### **Adults**

The adult dose varies from 25 mg (5 ml) to 400 mg (80 ml) spironolactone a day, depending on the condition being treated. If you are not sure how much to take, ask your doctor or pharmacist.

#### **Elderly**

Your doctor will start you on a low starting dose and gradually increase the dosage as needed to obtain the desired effect.

#### **Use in children and adolescents**

If you are giving Spironolactone Waymade to a child, the volume of this medicine you give will depend on the child's weight. Your doctor will work out the volume of this medicine that you should give.

#### **Method of administration**

An oral syringe and a bottle adapter are packed together with the oral liquid dosage form. The capacity of the syringe is 5 ml with scale graduation of 0.1 ml.

##### **Instruction for use**

1. Remove the bottle cap. Push the adapter into the opening of the bottle.
2. Insert the syringe into the hole in the adapter and turn the bottle upside down. Measure the dose by slowly pulling out the plunger to the correct volume. Read the dose on the top of the piston. Turn the bottle right and remove the syringe.
3. Discharge the syringe contents into the mouth.
4. Rinse the syringe and replace the cap.

#### **If you take more Spironolactone Waymade than you should**

If you accidentally take more Spironolactone Waymade, contact your doctor or nearest hospital accident and emergency department immediately. The symptoms of an overdose are feeling drowsy, dizzy, feeling dehydrated and you may feel confused. You may also feel or be sick, suffer from diarrhoea and may have skin rashes that will appear as flat red areas of skin with overlapping small raised bumps. Changes in your blood sodium and potassium levels may leave you feeling weak and suffering from tingling, prickling or numbness of the skin and/or muscle spasms but these symptoms are unlikely to be associated with severe overdosage.

**If you forget to take Spironolactone Waymade**

If you forget to take Spironolactone Waymade, take it as soon as you remember, unless it is almost time for your next dose. Do not take a double dose to make up for a forgotten dose.

**If you stop taking Spironolactone Waymade**

It is important to keep taking Spironolactone Waymade until your doctor tells you to stop, even if you start to feel better. If you stop taking Spironolactone Waymade too soon, your condition may get worse.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Tell your doctor immediately** if you experience any of the following symptoms after taking this medicine. Although they are very rare, the symptoms can be severe.

- Itchiness and blistering of the skin around the lips and the rest of the body, red or purple rash spreading and forming blisters (Stevens-Johnson syndrome)
- Detachment of the top layer of skin from the lower layers of skin, all over the body (toxic epidermal necrolysis - TEN)
- Skin rash, fever and swelling (which could be symptoms of something more serious, Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS))
- Yellow skin and eyes (jaundice)
- Spironolactone Waymade can cause impairment of liver function
- Irregular heartbeat which can be fatal, tingling sensation, paralysis (loss of muscle function) or difficulty in breathing; which may be symptoms of raised potassium levels in your blood. Your doctor will conduct regular blood tests to monitor potassium and other electrolyte levels. They may stop your treatment if necessary.

**List of other side effects of Spironolactone Waymade by frequency:**

Very common: may affect more than 1 in 10 people

- Raised potassium in the blood

Common: may affect up to 1 in 10 people

- Confusion
- Dizziness
- Vomiting or feeling sick
- Itching of the skin
- Rash
- Muscle or leg cramps
- Kidney failure or abnormal function
- Breast enlargement in men
- Breast pain (in men)
- Feeling generally unwell

Uncommon: may affect up to 1 in 100 people

- Changes in the breast such as breast lumps
- Disturbances in body electrolytes such as high blood calcium
- Abnormal functioning of the liver
- Skin allergy with development of itchiness and hives, nettle like rash
- Menstrual problems in women
- Breast pain (in women)

Not known: frequency cannot be estimated from the available data

- Lowered white blood cell count in blood
- Reduced number of cells that fight infection – white blood cells which make infections more likely
- Reduced number of cells that help with blood clotting which increases risk of bleeding or bruising
- Change in sex drive for both men and women
- Digestion problems, stomach upset
- Skin condition presenting with fluid-filled blisters (pemphigoid)
- Hair loss
- Excessive hair growth

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via Yellow Card Scheme website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Spironolactone Waymade**

Keep this medicine out of the sight and reach of children.

This medicinal product does not require any special temperature storage conditions. Store in the original package in order to protect from light.

Do not use this medicine after the expiry date which is stated on the bottle label or carton after EXP. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Spironolactone Waymade contains**

- The active substance is spironolactone.
- The other ingredients are Butylated hydroxyanisole (E 320), Peppermint oil, and Medium-chain triglycerides.

### **What Spironolactone Waymade looks like and contents of the pack**

Spironolactone Waymade is a clear liquid solution.

Spironolactone Waymade comes in amber glass bottle with child resistant cap. An oral syringe and a bottle adapter are packed together with the bottle. The capacity of the syringe is 5 ml with scale graduation of 0.1 ml.

Pack size: 125 ml

## **Marketing Authorisation Holder and Manufacturer**

### **Marketing Authorisation Holder**

Waymade PLC  
Sovereign House,  
Miles Gray Road, Basildon,  
Essex, SS14 3FR, UK

**Manufacturer**

LABORATORIUM SANITATIS, S.L.

C/ Leonardo da Vinci, 11

(Parque Tecnológico de Álava),

Miñano, 01510 Alava

Spain

Waymade PLC

Sovereign House,

Miles Gray Road, Basildon,

Essex, SS14 3FR, UK

**This leaflet was last revised in 09/2024.**

To request a copy of this leaflet in Braille, large print or audio format, contact the licence holder at the above address or telephone: 01268 535200 (select option Medical Information) / e-mail: [medinfo@waymade.co.uk](mailto:medinfo@waymade.co.uk)