

Package leaflet: Information for the user

Cytisine 1.5 mg tablets

Cytisine

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Cytisine is and what it is used for
2. What you need to know before you use Cytisine
3. How to use Cytisine
4. Possible side effects
5. How to store Cytisine
6. Contents of the pack and other information

1. What Cytisine is and what it is used for

Smoking cessation and reduction of nicotine cravings in smokers who are willing to stop smoking. The treatment goal of Cytisine is the permanent cessation of the nicotine-containing products use.

The use of Cytisine allows for a gradual reduction of nicotine dependence by relieving withdrawal symptoms.

2. What you need to know before you use Cytisine

Do not use Cytisine:

- if you are allergic to cytisine or any of the other ingredients of this medicine (listed in section 6).
- if you have unstable angina,
- if you have a history of recent myocardial infarction,
- if you have clinically significant cardiac arrhythmias,
- if you had stroke recently,
- if you are pregnant or breast-feeding.

Warnings and precautions

Cytisine should be used with caution in case of ischemic heart disease, heart failure, hypertension, pheochromocytoma (a tumour of the adrenal gland), atherosclerosis (hardening of the arteries) and other peripheral vascular diseases, gastric and duodenal ulcer, gastroesophageal reflux disease, hyperthyroidism (overactive thyroid), diabetes, schizophrenia, kidney and liver failure.

Cytisine should be taken only by those with a serious intention of weaning off nicotine. The use of Cytisine and continuation of smoking could lead to aggravated side effects of nicotine.

Children and adolescents

Due to limited experience, the drug is not recommended for use in persons under 18 years of age.

Elderly population

Due to limited clinical experience, Cytisine is not recommended for use in elderly patients over 65 years of age.

Patients with kidney and liver impairment

There is no clinical experience of Cytisine in patients with kidney or liver impairment, therefore the medicine is not recommended for use in these patients.

Other medicines and Cytisine

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines. Do not take Cytisine with anti-tuberculosis drugs.

In some cases as a result of stopping smoking, with or without Cytisine an adjustment of the dose of other medicines may be necessary. This is especially important if you use other medicines which contain theophylline (to treat asthma), tacrine (for Alzheimer's disease), clozapine (for schizophrenia) and ropinirole (to treat Parkinson's disease). If you are not sure, talk to your doctor or pharmacist.

It is currently unknown whether Cytisine may reduce the effectiveness of systemically acting hormonal contraceptives. If you are using systemically acting hormonal contraceptives, you should add a second barrier method (e.g. condoms).

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

You must use effective methods of contraception if you are a woman of childbearing potential. Ask your doctor for advice.

Cytisine is contraindicated during pregnancy and breastfeeding.

Driving and using machines

Cytisine has no influence on the ability to drive and use machines.

Stopping smoking

The effects of changes in your body resulting from stopping smoking, with or without treatment with Cytisine, may alter the way other medicines act. Therefore, in some cases an adjustment of the dose may be necessary. See above under 'Other medicines and Cytisine for further details.

For some people, stopping smoking with or without treatment has been associated with an increased risk of experiencing changes in thinking or behaviour, feelings of depression and anxiety (rarely including suicidal ideation and suicide attempt) and can be associated with a worsening of psychiatric disorder. If you have a history of psychiatric disorder you should discuss this with your doctor.

3. How to use Cytisine

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

One package of Cytisine (100 tablets) is sufficient for a complete therapy. The duration of treatment is 25 days. Cytisine is for oral use and it should be taken with a suitable amount of water according to the following schedule.

Days of treatment	Recommended dosing	Maximum daily dose
From the 1st to the 3rd day	1 tablet every 2 hours	6 tablets
From the 4th to the 12th day	1 tablet every 2.5 hours	5 tablets
From the 13th to the 16th day	1 tablet every 3 hours	4 tablets
From the 17th to the 20th day	1 tablet every 5 hours	3 tablets

Keep this medicine out of the sight and reach of children.

Store below 25°C in the original package in order to protect from moisture and light.

Do not use this medicine after the expiry date which is stated on the package. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Cytisine contains

- The active substance is cytisine. One tablet contains 1.5 mg of cytisine.
- The other ingredients are: mannitol, microcrystalline cellulose, magnesium stearate, glycerol dibehenate, hypromellose.

What Cytisine looks like and contents of the pack

Cytisine is round, biconvex, white tablet with diameter 6 mm.

PVC/PCTFE/Aluminium blisters placed into cardboard box containing 100 tablets.

Marketing Authorisation Holder

Bonteque Consulting Ltd
29 Westcott Crescent, Hanwell
W7 1PL London, United Kingdom

Manufacturer

Aflofarm Farmacja Polska Sp. z o.o.
Szkolna 31
95-054 Ksawerów
Poland

Distributed By:
Consilient Health Ltd.
1 Church Road, Richmond upon Thames,
Surrey, TW9 2QE, United Kingdom

This leaflet was last revised in 10/2023.

<distributor's logo>