

## Package leaflet: Information for the patient

### Dabigatran etexilate 150 mg hard capsules

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Dabigatran etexilate is and what it is used for
2. What you need to know before you take Dabigatran etexilate
3. How to take Dabigatran etexilate
4. Possible side effects
5. How to store Dabigatran etexilate
6. Contents of the pack and other information

### 1. What Dabigatran etexilate is and what it is used for

These capsules contain the active substance dabigatran etexilate and belongs to a group of medicines called anticoagulants. It works by blocking a substance in the body which is involved in blood clot formation.

Dabigatran etexilate is used in adults to:

- prevent blood clots in the brain (stroke) and other blood vessels in the body if you have a form of irregular heart rhythm called nonvalvular atrial fibrillation and at least one additional risk factor.
- treat blood clots in the veins of your legs and lungs and to prevent blood clots from reoccurring in the vein of your legs and lungs.

Dabigatran etexilate is used in children to:

- treat blood clots and to prevent blood clots from reoccurring.

### 2. What you need to know before you take Dabigatran etexilate

#### Do not take Dabigatran etexilate if you;

- are allergic to dabigatran etexilate or any of the other ingredients of this medicine (listed in section 6).
- have severely reduced kidney function.
- are currently bleeding.
- have a disease in an organ of the body that increases the risk of serious bleeding (e.g., stomach ulcer, injury or bleeding in the brain, recent surgery of the brain or eyes).
- have an increased tendency to bleed. This may be inborn, of unknown cause or due to other medicines.
- are taking medicines to prevent blood clotting (e.g. warfarin, rivaroxaban, apixaban or heparin), except when changing anticoagulant treatment, while having a venous or arterial line and you get heparin through this line to keep it open or while your heart beat is being restored to normal by a procedure called catheter ablation for atrial fibrillation.
- have a severely reduced liver function or liver disease which could possibly cause death.
- are taking oral ketoconazole or itraconazole, medicines to treat fungal infections.
- are taking oral ciclosporine, a medicine to prevent organ rejection after transplantation.
- are taking dronedarone, a medicine used to treat abnormal heart beat.
- are taking a combination product of glecaprevir and pibrentasvir, an antiviral medicine used to treat hepatitis C.
- have received an artificial heart valve which requires permanent blood thinning.

#### Warnings and precautions

Talk to your doctor before taking Dabigatran etexilate. You may also need to talk to your doctor during treatment with this medicine if you experience symptoms or if you have to undergo surgery.

Tell your doctor if you have or have had any medical conditions or illnesses, in particular any of those included in the following list:

- if you have an increased bleeding risk, such as:
  - if you have been recently bleeding.
  - if you have had a surgical tissue removal (biopsy) in the past month.
  - if you have had a serious injury (e.g. a bone fracture, head injury or any injury requiring surgical treatment).
  - if you are suffering from an inflammation of the gullet or stomach.
  - if you have problems with reflux of gastric juice into the gullet.
- if you are receiving medicines which could increase the risk of bleeding. See 'Other medicines and Dabigatran etexilate' below.
- if you are taking anti-inflammatory medicines such as diclofenac, ibuprofen, piroxicam.
- if you are suffering from an infection of the heart (bacterial endocarditis).
- if you know you have impaired kidney function, or you are suffering from dehydration (symptoms include feeling thirsty and passing reduced amounts of dark-coloured (concentrated) urine).
- if you are older than 75 years.
- if you are an adult patient and weigh 50 kg or less.
- only if used for children: if the child has an infection around or within the brain.
- if you have had a heart attack or if you have been diagnosed with conditions that increase the risk to develop a heart attack.
- if you have a liver disease that is associated with changes in the blood tests. The use of Dabigatran etexilate is not recommended in this case.

#### Take special care with Dabigatran etexilate

- if you need to have an operation:

In this case Dabigatran etexilate will need to be stopped temporarily due to an increased bleeding risk during and shortly after an operation. It is very important to take Dabigatran etexilate before and after the operation exactly at the times you have been told by your doctor.
- if an operation involves a catheter or injection into your spinal column (e.g. for epidural or spinal anaesthesia or pain reduction):
  - it is very important to take Dabigatran etexilate before and after the operation exactly at the times you have been told by your doctor.
  - tell your doctor immediately if you get numbness or weakness of your legs or problems with your bowel or bladder after the end of anaesthesia, because urgent care is necessary.
- if you fall or injure yourself during treatment, especially if you hit your head. Please seek urgent medical attention. You may need to be checked by a doctor, as you may be at increased risk of bleeding.
- if you know that you have a disease called antiphospholipid syndrome (a disorder of the immune system that causes an increased risk of blood clots), tell your doctor who will decide if the treatment may need to be changed.

#### Other medicines and Dabigatran etexilate

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. **In particular you should tell your doctor before taking Dabigatran etexilate, if you are taking one of the medicines listed below:**

- Medicines to reduce blood clotting (e.g. warfarin, phenprocoumon, acenocoumarol, heparin, clopidogrel, prasugrel, ticagrelor, rivaroxaban, acetylsalicylic acid).
- Medicines to treat fungal infections (e.g. ketoconazole, itraconazole), unless they are only applied to the skin.
- Medicines to treat abnormal heart beats (e.g. amiodarone, dronedarone, quinidine, verapamil). If you are taking verapamil containing medicines, your doctor may tell you to use a reduced dose of Dabigatran etexilate depending on the condition for which it is prescribed to you. See section 3.
- Medicines to prevent organ rejection after transplantation (e.g. tacrolimus, ciclosporine).
- A combination product of glecaprevir and pibrentasvir (an antiviral medicine used to treat hepatitis C).
- Anti-inflammatory and pain reliever medicines (e.g. acetylsalicylic acid, ibuprofen, diclofenac).
- St. John's wort, a herbal medicine for depression.
- Antidepressant medicines called selective serotonin re-uptake inhibitors or serotonin- norepinephrine re-uptake inhibitors.
- Rifampicin or clarithromycin (two antibiotics).
- Anti-viral medicines for AIDS (e.g. ritonavir).
- Certain medicines for treatment of epilepsy (e.g. carbamazepine, phenytoin).

#### Pregnancy and breast-feeding

The effects of Dabigatran etexilate on pregnancy and the unborn child are not known. You should not take this medicine if you are pregnant unless your doctor advises you that it is safe to do so. If you are a woman of child-bearing age, you should avoid becoming pregnant while you are taking Dabigatran etexilate.

You should not breast-feed while you are taking Dabigatran etexilate.

#### Driving and using machines

Dabigatran etexilate has no known effects on the ability to drive or use machines.

### 3. How to take Dabigatran etexilate

Dabigatran etexilate capsules can be used in adults and children aged 8 years or older who are able to swallow the capsules whole. There are other age appropriate dose forms for the treatment of children below 8 years.

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

#### Take Dabigatran etexilate as recommended for the following conditions:

**Prevention of brain or body vessel obstruction by blood clot formation developing after abnormal heart beats and treatment of blood clots in the veins of your legs and lungs including prevention of blood clots from re-occurring in the vein of your legs and lungs**

The recommended dose is 300 mg taken as one **150 mg capsule twice a day**.

If you are **80 years or older**, the recommended dose of Dabigatran etexilate is 220 mg taken as **one 110 mg capsule twice daily**.

If you are taking **verapamil containing medicines**, you should be treated with a reduced Dabigatran etexilate dose of 220 mg taken as **one 110 mg capsule twice a day**, because your bleeding risk may be increased.

If you have a **potentially higher risk for bleeding**, your doctor may decide to prescribe a dose of Dabigatran etexilate 220 mg taken as **one 110 mg capsule twice a day**.

You can continue to take this medicine if your heart beat needs to be restored to normal by a procedure called cardioversion or by a procedure called catheter ablation for atrial fibrillation. Take Dabigatran etexilate as your physician has told you.

If a medical device (stent) has been deployed in a blood vessel to keep it open in a procedure called percutaneous coronary intervention with stenting, you can be treated with Dabigatran etexilate after your physician has decided that normal control of blood coagulation is achieved. Take Dabigatran etexilate as your physician has told you.

#### Treatment of blood clots and prevention of blood clots from reoccurring in children

**Dabigatran etexilate should be taken twice daily**, one dose in the morning and one dose in the evening, at approximately the same time every day. The dosing interval should be as close to 12 hours as possible.

The recommended dose depends on weight and age. Your doctor will determine the correct dose. Your doctor may adjust the dose as treatment progresses. Keep using all other medicines, unless your doctor tells you to stop using any.

Table 1 shows single and total daily Dabigatran etexilate doses in milligrams (mg). The doses depend on weight in kilograms (kg) and age in years of the patient.

Table 1: Dosing table for Dabigatran etexilate capsules

Weight/age combinations		Single dose in mg	Total daily dose in mg
Weight in kg	Age in years		
11 to less than 13 kg	8 to less than 9 years	75	150
13 to less than 16 kg	8 to less than 11 years	110	220
16 to less than 21 kg	8 to less than 14 years	110	220
21 to less than 26 kg	8 to less than 16 years	150	300
26 to less than 31 kg	8 to less than 18 years	150	300
31 to less than 41 kg	8 to less than 18 years	185	370
41 to less than 51 kg	8 to less than 18 years	220	440
51 to less than 61 kg	8 to less than 18 years	260	520
61 to less than 71 kg	8 to less than 18 years	300	600
71 to less than 81 kg	8 to less than 18 years	300	600
81 kg or greater	10 to less than 18 years	300	600

Single doses requiring combinations of more than one capsule:

300 mg: two 150 mg capsules or four 75 mg capsules  
260 mg: one 110 mg plus one 150 mg capsule or one 110 mg plus two 75 mg capsules

220 mg: two 110 mg capsules

185 mg: one 75 mg plus one 110 mg capsule

150 mg: one 150 mg capsule or two 75 mg capsules

#### How to take Dabigatran etexilate

Dabigatran etexilate can be taken with or without food. The capsule should be swallowed whole with a glass of water, to ensure delivery to the stomach. Do not break, chew, or empty the pellets from the capsule since this may increase the risk of bleeding.

#### Instructions for opening the blisters

- Tear off one individual blister from the blister card along the perforated line
- Peel off the backing foil and remove the capsule.
- Do not push the capsules through the blister foil.
- Do not peel off the blister foil until a capsule is required.

#### Change of anticoagulant treatment

Without specific guidance from your doctor do not change your anticoagulant treatment.

#### If you take more Dabigatran etexilate than you should

Taking too much of this medicine increases the risk of bleeding. Contact your doctor immediately if you have taken too many Dabigatran etexilate capsules. Specific treatment options are available.

#### If you forget to take Dabigatran etexilate

A forgotten dose can still be taken up to 6 hours prior to the next due dose.

A missed dose should be omitted if the remaining time is below 6 hours prior to the next due dose. Do not take a double dose to make up for a forgotten dose.

#### If you stop taking Dabigatran etexilate

Take Dabigatran etexilate exactly as prescribed. Do not stop taking Dabigatran etexilate without talking to your doctor first, because the risk of developing a blood clot could be higher if you stop treatment too early. Contact your doctor if you experience indigestion after taking Dabigatran etexilate.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.



#### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Dabigatran etexilate affects blood clotting, so most side effects are related to signs such as bruising or bleeding.

Major or severe bleeding may occur, these constitute the most serious side effects and, regardless of location, may become disabling, life-threatening or even lead to death. In some cases these bleedings may not be obvious.

If you experience any bleeding event that does not stop by itself or if you experience signs of excessive bleeding (exceptional weakness, tiredness, paleness, dizziness, headache or unexplained swelling) consult your doctor immediately. Your doctor may decide to keep you under closer observation or change your medicine.

Tell your doctor immediately, if you experience a serious allergic reaction which causes difficulty in breathing or dizziness.

Possible side effects are listed below, grouped by how likely they are to happen.

##### Prevention of brain or body vessel obstruction by blood clot formation developing after abnormal heart beats

**Common** (may affect up to 1 in 10 people):

- Bleeding may happen from the nose, into the stomach or bowel, from penis/vagina or urinary tract (incl. blood in the urine that stains the urine pink or red), or under the skin.
- A fall in the number of red cells in the blood.
- Belly ache or stomach ache.
- Indigestion.
- Frequent loose or liquid bowel movements.
- Feeling sick.

**Uncommon** (may affect up to 1 in 100 people):

- Bleeding.
- Bleeding may happen from piles, from the rectum, or in the brain.
- Haematoma formation.
- Coughing of blood or blood stained sputum.
- A fall in the number of platelets in the blood.
- A fall in the amount of haemoglobin in the blood (the substance in the red blood cells).
- Allergic reaction.
- Sudden change of the skin which affects its colour and appearance.
- Itching.
- Ulcer in the stomach or bowel (incl. ulcer in the gullet).
- Inflammation of the gullet and stomach.
- Reflux of gastric juice into the gullet.
- Vomiting.
- Difficulty in swallowing.
- Unusual laboratory test results on liver function.

**Rare** (may affect up to 1 in 1,000 people):

- Bleeding may happen into a joint, from a surgical incision, from an injury, from the site of entry of an injection or from the site of entry of a catheter into a vein.
- Serious allergic reaction which causes difficulty in breathing or dizziness.
- Serious allergic reaction which causes swelling of the face or throat.
- Skin rash notable for dark red, raised, itchy bumps caused by an allergic reaction.
- A decrease in the proportion of blood cells.
- Liver enzymes increased.
- Yellowing of the skin or whites of the eyes, caused by liver or blood problems.

**Not known** (frequency cannot be estimated from the available data):

- Difficulty in breathing or wheezing.
- Decreases in the number or even lack of white blood cells (which help to fight infections).
- Hair loss.

In a clinical trial the rate of heart attacks with Dabigatran etexilate was numerically higher than with warfarin. The overall occurrence was low.

##### Treatment of blood clots in the veins of your legs and lungs including prevention of blood clots from re-occurring in the veins of your legs and/or lungs

**Common** (may affect up to 1 in 10 people):

- Bleeding may happen from the nose, into the stomach or bowel, from the rectum, from penis/vagina or urinary tract (incl. blood in the urine that stains the urine pink or red), or under the skin.
- Indigestion.

**Uncommon** (may affect up to 1 in 100 people):

- Bleeding.
- Bleeding may happen into a joint or from an injury.
- Bleeding may happen from piles.
- A fall in the number of red cells in the blood.
- Haematoma formation.
- Coughing of blood or blood stained sputum.
- Allergic reaction.
- Sudden change of the skin which affects its colour and appearance.
- Itching.
- Ulcer in the stomach or bowel.
- Inflammation of the gullet and stomach.
- Reflux of gastric juice into the gullet.
- Feeling sick.
- Vomiting.
- Belly ache or stomach ache.
- Frequent loose or liquid bowel movements.
- Unusual laboratory test results on liver function.
- Liver enzymes increased.

**Rare** (may affect up to 1 in 1,000 people):

- Bleeding may happen, from a surgical incision, or from the site of entry of an injection or from the site of entry of a catheter into a vein or from the brain.
- A fall in the number of platelets in the blood.
- Serious allergic reaction which causes difficulty in breathing or dizziness.
- Serious allergic reaction which causes swelling of the face or throat.
- Skin rash notable for dark red, raised, itchy bumps caused by an allergic reaction.
- Difficulty in swallowing.
- A decrease in the proportion of red cells in the blood.

**Not known** (frequency cannot be estimated from the available data):

- Difficulty in breathing or wheezing.
- A fall in the amount of haemoglobin in the blood (the substance in the red blood cells).
- A fall in the number of red cells in the blood.
- Decrease in the number or even lack of white blood cells (which help to fight infections).
- Yellowing of the skin or whites of the eyes, caused by liver or blood problems.
- Hair loss.

In the trial program the rate of heart attacks with Dabigatran etexilate was higher than with warfarin. The overall occurrence was low. No imbalance in the rate of heart attacks was observed in patients treated with dabigatran versus patients treated with placebo.

##### Treatment of blood clots and prevention of blood clots from reoccurring in children

**Common** (may affect up to 1 in 10 people):

- A fall in the number of red cells in the blood.
- A fall in the number of platelets in the blood.
- Skin rash notable for dark red, raised, itchy bumps caused by an allergic reaction.
- Sudden change of the skin which affects its colour and appearance.
- Haematoma formation.
- Nosebleed.
- Reflux of gastric juices into the gullet.
- Vomiting.
- Feeling sick.
- Frequent loose or liquid bowel movements.
- Indigestion.
- Hair loss.
- Liver enzymes increased.

**Uncommon** (may affect up to 1 in 100 people):

- Decrease in the number of white blood cells (which help to fight infections)
- Bleeding may happen into the stomach or bowel, from the brain, from the rectum, from penis/vagina or urinary tract (incl. blood in the urine that stains the urine pink or red), or under the skin.
- A fall in the amount of haemoglobin in the blood (the substance in the red blood cells).
- A decrease in the proportion of blood cells.
- Itching.
- Coughing of blood or blood stained sputum.
- Belly ache or stomach ache.
- Inflammation of the gullet and stomach.
- Allergic reaction.
- Difficulty in swallowing.
- Yellowing of the skin or whites of the eyes, caused by liver or blood problems.

**Not known** (frequency cannot be estimated from the available data):

- Lack of white blood cells (which help to fight infections).
- Serious allergic reaction which causes difficulty in breathing or dizziness.
- Serious allergic reaction which causes swelling of the face or throat.
- Difficulty in breathing or wheezing.
- Bleeding.
- Bleeding may happen into a joint or from an injury, from a surgical incision, or from the site of entry of an injection or from the site of entry of a catheter into a vein.
- Bleeding may happen from piles.
- Ulcer in the stomach or bowel (incl. ulcer in the gullet).
- Unusual laboratory test results on liver function.

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

#### 5. How to store Dabigatran etexilate

Keep this medicine out of the sight and reach of children. Do not use this medicine after the expiry date which is stated on the carton or blister after “EXP”. The expiry date refers to the last day of that month.

Store in the original package in order to protect from moisture.

Do not throw away any medicines via wastewater. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

#### 6. Contents of the pack and other information

**What Dabigatran etexilate contains**

- The active substance is dabigatran etexilate. Each hard capsule contains 150 mg dabigatran etexilate (as mesilate).
- The other ingredients are tartaric acid, acacia, hypromellose, dimeticone, talc and hydroxypropylcellulose.
- The capsule shell contains carrageenan, potassium chloride, titanium dioxide (E-171), hypromellose and indigo carmine (E-132).

**What Dabigatran etexilate looks like and contents of the pack**

Dabigatran etexilate 150 mg are hard capsules containing off white to pale yellow coloured pellets filled in blue cap and white to off white body hard capsules size 0.

Dabigatran etexilate is available in packs containing 1 x 10, 3 x 10 or 6 x 10 hard capsules in perforated Aluminum/OPA- ALU-PVC blisters.

Not all pack sizes may be marketed.

**Marketing Authorisation Holder:**

Zentiva Pharma UK Limited,  
12 New Fetter Lane,  
London, EC4A 1JP,  
United Kingdom

**Manufacturers:**

Galenicum Health, S.L.  
Avda. Cornellá 144, 7<sup>o</sup>-1<sup>a</sup> Edificio Lekla,  
Esplugues de Llobregat,  
08950 Barcelona,  
Spain

or

SAG Manufacturing S.L.U.,  
Crta. N-I, Km 36,  
San Agustín de Guadalix,  
28750 Madrid,  
Spain

**This leaflet was last revised in April 2024**

#### Other sources of information

For information on large print, Braille or audio CD call emc accessibility on 0800 198 5000