

85.6 mm

53.98 mm



Fingolimod

PREGNANCY-SPECIFIC Patient Reminder Card

Date of preparation: April 2023
Approved by MHRA 05/2023

Version 1.0

Page 1

Before starting Fingolimod treatment

Fingolimod is contraindicated in pregnant women and women of child-bearing potential (including adolescents) not using effective contraception.

At treatment start and then regularly, your doctor will inform you about the teratogenic risk (causes defects to unborn babies) and required actions to minimise this risk.

Page 2

A pregnancy test must be conducted and the negative result verified by a doctor before starting treatment.

Your doctor will inform you about the need for effective contraception while on treatment and for 2 months after discontinuation. Talk to your doctor about the most effective contraception options available to you.

Please read the fingolimod patient guide leaflet provided by your doctor.

Page 3

While you are taking fingolimod

While on treatment women must not become pregnant.

Patients must use effective contraception while taking fingolimod.

Women must not become pregnant during treatment and for 2 months after discontinuing treatment.

Pregnancy tests must be repeated at suitable intervals. Your doctor will provide regular counselling about fingolimod's serious risk to the foetus.

Page 4

If you become pregnant or if you want to become pregnant, please discuss this with your doctor because fingolimod treatment must be discontinued.

In the event of a pregnancy your doctor will provide counselling.

Your doctor will give you medical advice regarding the harmful effects of fingolimod to the foetus and will provide an evaluation of the potential outcome.

Page 5

Dimensions: 85.6 mm x 53.98 mm



Key Lines not to be Printed



Pantone 273 C



Pantone 485 C



Black

After stopping fingolimod treatment

Inform your doctor immediately if you believe your MS is getting worse (e.g. weakness or visual changes) or if you notice any new symptoms after stopping treatment with fingolimod due to pregnancy

Effective contraception is needed for 2 months after stopping fingolimod treatment because of the length of time it takes for fingolimod to leave the body.

Page 6

Reporting of side effects

Please report suspected adverse drug reactions (ADRs) to the MHRA through the Yellow Card scheme, via the Yellow Card website www.mhra.gov.uk/yellowcard, the free Yellow Card app available in Apple App Store or Google Play Store, and also some clinical IT systems for healthcare professionals.

Alternatively, you can call 0800 731 6789 for free, Monday to Friday between 9 am and 5 pm.

Page 7

By reporting side effects, you can help provide more information on the safety of this medicine.

All pregnancies and adverse events should be reported to MSN Laboratories Europe Limited via email to pharmacovigilance@msnlabs.com or by calling 020 3371 7158.

Page 8

For further information please also contact MSN Laboratories Europe Limited via email to pharmacovigilance@msnlabs.com or by calling 020 3371 7158.

Page 9