


Serialization	
PC	00000000000000
SN	00000000000000
Lot	00000000
EXP	00.0000



Preprinted on the box
Inkjet on the packaging line

- Cut
- Fold
- Braille
- Info (non printing ink)



PC
SN
Lot
EXP

Print/varnish free area

Please apply pharmacy label here

Apixaban 2.5 mg film-coated tablets
Each film-coated tablet contains 2.5 mg apixaban. Also contains lactose. See leaflet for further information. Oral use. Read the package leaflet before use. **Keep out of the sight and reach of children.**

Print/varnish free area

Apixaban 2.5 mg film-coated tablets
60 film-coated tablets

941252

Apixaban 2.5 mg film-coated tablets

8 594739 310162

13395

Apixaban 2.5 mg film-coated tablets
60 film-coated tablets

PLEASE FLIP UP THE PATIENT ALERT CARD AND ATTACH THE DISPENSING LABEL UNDERNEATH.

ZENTIVA

UK Only


Apixaban 2.5 mg film-coated tablets

PL 17780/1187 POM


Zentiva Pharma UK Limited,
12 New Fetter Lane, London, EC4A 1JP,
United Kingdom

<p>APIXABAN PATIENT ALERT CARD</p> <p>Last updated: 03/2024</p> <p>Carry this card with you at all times.</p> <p>Show this card to your pharmacist, dentist and any other healthcare professionals that treat you.</p> <p>I am under anticoagulation treatment with Apixaban to prevent blood clots.</p> <p>ZENTIVA</p>	<p>Please complete this section or ask your doctor to do it.</p> <p>Name: _____</p> <p>Date of Birth: _____</p> <p>Indication: _____</p> <p>Dose: _____ mg twice daily</p> <p>Doctor's Name: _____</p> <p>Doctor's telephone: _____</p>
--	--

Special glue



Print/varnish free area



<p>Information for patients</p> <ul style="list-style-type: none">• Take Apixaban regularly as instructed. If you miss a dose, take it as soon as you remember and continue to follow your dosing schedule.• Do not stop taking Apixaban without talking to your doctor, as you are at risk of suffering from a stroke or other complications.• Apixaban helps to thin your blood. However, this may increase your risk of bleeding.	<ul style="list-style-type: none">• Signs and symptoms of bleeding include bruising or bleeding under the skin, tar-coloured stools, blood in urine, nose-bleed, dizziness, tiredness, paleness or weakness, sudden severe headache, coughing up blood or vomiting blood.• If the bleeding does not stop on its own, seek medical attention immediately.• If you need surgery or any invasive procedure, inform your doctor that you are taking Apixaban.
<p>Information for healthcare professionals</p> <ul style="list-style-type: none">• Apixaban is an oral anticoagulant acting by direct selective inhibition of factor Xa.• Apixaban may increase the risk of bleeding. In case of major bleeding events, it should be stopped immediately.• Treatment with Apixaban does not require routine monitoring of exposure. A calibrated quantitative anti-Factor Xa assay may be useful in exceptional	<p>situations, e.g., overdose and emergency surgery (prothrombin time (PT), international normalised ratio (INR) and activated partial thromboplastin time (aPTT) clotting tests are not recommended) – see SmPC.</p> <ul style="list-style-type: none">• An agent to reverse the anti-factor Xa activity of apixaban is available.